



2.1(g) All strings over $\{a, b\}$ in which both the number of a 's and the number of b 's is even.



2.1(g2) All strings over $\{a, b\}$ in which either the number of a 's or the number of b 's is odd (or both).




Definition (FA)

[deterministic] finite automaton 5-tuple $M = (Q, \Sigma, q_0, A, \delta)$,


- Q finite set *states*;
- Σ finite *input alphabet*;
- $q_0 \in Q$ *initial state*;
- $A \subseteq Q$ *accepting states*;
- $\delta : Q \times \Sigma \rightarrow Q$ *transition function*.

[M] D 2.11 Finite automaton

[L] D 2.1 Deterministic finite accepter, has 'final' states



Example

$$L_1 = \{ x \in \{a, b\}^* \mid x \text{ ends with } aa \}$$


δ	a	b
q_0	q_1	q_0
q_1	q_2	q_0
q_2	q_2	q_0

[M] E. 2.1

Definition (FA)

[deterministic] finite automaton 5-tuple $M = (Q, \Sigma, q_0, A, \delta)$,

- Q finite set *states*;
- Σ finite *input alphabet*;
- $q_0 \in Q$ *initial state*;
- $A \subseteq Q$ *accepting states*;
- $\delta : Q \times \Sigma \rightarrow Q$ *transition function*.

[M] D 2.11 Finite automaton

[L] D 2.1 Deterministic finite accepter, has 'final' states

FA $M = (Q, \Sigma, q_0, A, \delta)$

Definition

extended transition function $\delta^* : Q \times \Sigma^* \rightarrow Q$, such that

- $\delta^*(q, \Lambda) = q$ for $q \in Q$
- $\delta^*(q, y\sigma) = \delta(\delta^*(q, y), \sigma)$ for $q \in Q, y \in \Sigma^*, \sigma \in \Sigma$



[M] D 2.12 [L] p.40/1

Theorem

$q = \delta^*(p, w)$ iff there is a path in [the transition graph of] M from p to q with label w .

[L] Th 2.1

Extended transition function



$$\delta^*(q_0, aabb) = q_0 :$$

$$\delta^*(q_0, \Lambda) = q_0$$

$$\delta^*(q_0, a) = \delta^*(q_0, \Lambda a) = \delta(\delta^*(q_0, \Lambda), a) = \delta(q_0, a) = q_1$$

$$\delta^*(q_0, aa) = \delta(\delta^*(q_0, a), a) = \delta(q_1, a) = q_2$$

$$\delta^*(q_0, aab) = \delta(\delta^*(q_0, aa), b) = \delta(q_2, b) = q_0$$

$$\delta^*(q_0, aabb) = \delta(\delta^*(q_0, aab), b) = \delta(q_0, b) = q_0$$

Definition


Let $M = (Q, \Sigma, q_0, A, \delta)$ be an FA, and let $x \in \Sigma^*$. The string x is *accepted* by M if $\delta^*(q_0, x) \in A$.

The *language accepted* by $M = (Q, \Sigma, q_0, A, \delta)$ is the set $L(M) = \{ x \in \Sigma^* \mid x \text{ is accepted by } M \}$

[M] D 2.14 [L] D 2.2

From lecture 1:

$$L_2 = \{ x \in \{a, b\}^* \mid x \text{ ends with } b \text{ and does not contain } aa \}$$



$$\neg(P \wedge Q) = \neg P \vee \neg Q$$

$$L_2^c = \{ x \in \{a, b\}^* \mid x \text{ does not end with } b \text{ or contains } aa \}$$

Construction




FA $M = (Q, \Sigma, q_0, A, \delta)$,

let $M^c = (Q, \Sigma, q_0, Q - A, \delta)$


Theorem

$$L(M^c) = \Sigma^* - L(M)$$

Proof...

Example (Even number of a , and ending with b)

Even number of *a* and ending with *b*



FA $M_i = (Q_i, \Sigma, q_i, A_i, \delta_i) \quad i = 1, 2$

Product construction

construct FA $M = (Q, \Sigma, q_0, A, \delta)$ such that

- $Q = Q_1 \times Q_2$
- $q_0 = (q_1, q_2)$
- $\delta((p, q), \sigma) = (\delta_1(p, \sigma), \delta_2(q, \sigma))$
- A as needed

Theorem (2.15 Parallel simulation)

- $A = \{(p, q) \mid p \in A_1 \text{ or } q \in A_2\}$, then $L(M) = L(M_1) \cup L(M_2)$
- $A = \{(p, q) \mid p \in A_1 \text{ and } q \in A_2\}$, then $L(M) = L(M_1) \cap L(M_2)$
- $A = \{(p, q) \mid p \in A_1 \text{ and } q \notin A_2\}$, then $L(M) = L(M_1) - L(M_2)$

Proof...

[M] Sect 2.2



Exercise 2.11.

Use induction to show that for every $x \in \Sigma^*$ and every $(p, q) \in Q$,

$$\delta^*((p, q), x) = (\delta_1^*(p, x), \delta_2^*(q, x))$$

Example: intersection 'and' (product construction)

not substring aa





ends with ab


[M] E 2.16

Example: intersection 'and' (product construction)

not substring aa




ends with ab





[M] E 2.16


Example: union, contain either ab or bba



[M] E. 2.18, see also \hookrightarrow subset construction

$L = \{ w \in \{a, b\}^* \mid w \text{ begint en eindigt met een } a, \text{ en } |w| \text{ is even} \}$





[M] Fig. 2.28

Pumping lemma for regular languages

Regular language is language accepted by an FA.

Theorem

Suppose L is a language over the alphabet Σ . If L is accepted by a finite automaton M , and if n is the number of states of M , then

- \forall for every $x \in L$
satisfying $|x| \geq n$
- \exists there are three strings u , v , and w ,
such that $x = uvw$ and the following three conditions are true:
 - (1) $|uv| \leq n$,
 - (2) $|v| \geq 1$
- \forall and (3) for all $m \geq 0$, $uv^m w$ belongs to L

[M] Thm. 2.29

Pumping lemma for regular languages

In other words:

Theorem

- ∀ For every regular language L
- ∃ there exists a constant $n \geq 1$
such that
- ∀ for every $x \in L$
with $|x| \geq n$
- ∃ there exists a decomposition $x = uvw$
with (1) $|uv| \leq n$,
and (2) $|v| \geq 1$
such that
- ∀ (3) for all $m \geq 0$, $uv^m w \in L$

if $L = L(M)$ then $n = |Q|$.

[M] Thm. 2.29

Pumping lemma for regular languages

In other words:

Theorem

If L is a regular language, then

\exists *there exists a constant $n \geq 1$
such that*

\forall *for every $x \in L$
with $|x| \geq n$*

\exists *there exists a decomposition $x = uvw$
with (1) $|uv| \leq n$,
and (2) $|v| \geq 1$
such that*

\forall *(3) for all $m \geq 0$, $uv^m w \in L$*

if $L = L(M)$ then $n = |Q|$.

Introduction to Logic: $p \rightarrow q \iff \neg q \rightarrow \neg p$

Theorem

If

\forall for every $n \geq 1$

\exists there exists $x \in L$
with $|x| \geq n$
such that

\forall for every decomposition $x = uvw$
with (1) $|uv| \leq n$,
and (2) $|v| \geq 1$

\exists (3) there exists $m \geq 0$,
such that
 $uv^m w \notin L$

then L is not a regular language.

[M] Thm. 2.29

Example

$L = \{a^i b^i \mid i \geq 0\}$ is not accepted by FA.

[M] E 2.30

Proof: by contradiction

We prove that the language $L = \{a^i b^i \mid i \geq 0\}$ is not regular, by contradiction.

Assume that $L = \{a^i b^i \mid i \geq 0\}$ is accepted by FA with n states.

Take $x = a^n b^n$. Then $x \in L$, and $|x| = 2n \geq n$.

Thus there exists a decomposition $x = uvw$ such that $|uv| \leq n$ with v nonempty, and $uv^m w \in L$ for every m .

Whatever this decomposition is, v consists of a 's only. Consider $m = 0$. Deleting v from the string x will delete a number of a 's. So $uv^0 w$ is of the form $a^{n'} b^n$ with $n' < n$.

This string is not in L ; a contradiction. ($m \geq 2$ would also yield contradiction)

So, L is not regular.

Example

$L = \{a^i b^i \mid i \geq 0\}$ is not accepted by FA.

[M] E 2.30

$AeqB = \{x \in \{a, b\}^* \mid n_a(x) = n_b(x)\}$

Same argument, or closure properties

FA $M_i = (Q_i, \Sigma, q_i, A_i, \delta_i) \quad i = 1, 2$

Product construction

construct FA $M = (Q, \Sigma, q_0, A, \delta)$ such that

- $Q = Q_1 \times Q_2$
- $q_0 = (q_1, q_2)$
- $\delta((p, q), \sigma) = (\delta_1(p, \sigma), \delta_2(q, \sigma))$
- A as needed

Theorem (2.15 Parallel simulation)

- $A = \{(p, q) \mid p \in A_1 \text{ or } q \in A_2\}$, then $L(M) = L(M_1) \cup L(M_2)$
- $A = \{(p, q) \mid p \in A_1 \text{ and } q \in A_2\}$, then $L(M) = L(M_1) \cap L(M_2)$
- $A = \{(p, q) \mid p \in A_1 \text{ and } q \notin A_2\}$, then $L(M) = L(M_1) - L(M_2)$

Proof...

[M] Sect 2.2

Exactly the same argument can be used (verbatim) to prove that $L = \text{AeqB}$ is not regular.

We can also apply closure properties of REG to see that AeqB is not regular, as follows.

Assume AeqB is regular. Then also $\text{AnBn} = \text{AeqB} \cap \mathbf{a^*b^*}$ is regular, as regular languages are closed under intersection.

This is a contradiction, as we just have argued that AnBn is not regular. Thus, also AeqB is not regular.

Issues:

- Which n ? Can I just take $x = aababaabbab$?
- Which x ? Some x may not yield a contradiction.
- Which decomposition uvw ? Can I just take $u = a^{10}$, $v = a^{n-10}$, $w = b^n$?
- Which m ? Some m may not yield a contradiction.

Example

$L = \{ x \in \{a, b\}^* \mid n_a(x) > n_b(x) \}$ is not accepted by FA

[M] E 2.31